

Final Draft
of the original manuscript:

Jung, F.; Rampling, M.:
Editorial: **Role of blood viscosity in the microcirculation**
In: Clinical Hemorheology and Microcirculation (2016) IOS
Press

DOI: 10.3233/CH-168108

Role of blood viscosity in the microcirculation

F. Jung^a and M. Rampling^b

^a Institute of Biomaterial Science and Berlin-Brandenburg Centre for Regenerative Therapies, Helmholtz-Zentrum Geesthacht, Teltow, Germany

^b Department of Bioengineering, Imperial College, London SW7 2AZ, UK.

Blood fluidity is reported to influence capillary perfusion in healthy subjects and especially in patients with different diseases [6, 10, 29, 31]. Up to now there is no simple way to quantify the influence of the different haemorheological parameters (haematocrit, plasma viscosity, erythrocyte aggregation, erythrocyte deformability) on the velocity of blood cells in human capillaries. In general, only a limited correlation can be shown [18]. One reason is the great variability of the diameter of human capillaries described in healthy subjects, with values between 3 and 15 μm , and in patients - depending on the disease – with values up to 50 μm (e.g. giant capillaries in patients with scleroderma [33]). This is a very large range with respect to the role of the various haemorheological parameters of blood in microcirculation.

The hypothesis of Barras, that plasma viscosity determines the perfusion of microvessels because the absolute capillary haematocrit is about 10 - 20%, so that blood viscosity in capillaries approaches plasma viscosity seems to be near the truth [4], as long as the capillary diameter is big in relation to the erythrocyte diameter. The work of Lipowsky would seem to confirm this [25]. In the case where the capillary diameter is smaller than the red blood cell diameter, the cells have to be deformed by shear forces before passing through a capillary. In this case the blood flow velocity should additionally depend on the erythrocyte deformability. This fact, which results from a fluid-dynamical point of view, however, is often not seen in patients using capillary microscopy in nailfold capillaries: neighbouring capillaries with different diameters and fed from the same arteriole show the same red blood cell velocity. Therefore, we conclude that there is a regulatory mechanism in the capillary network possibly by the microvascular endothelium as described by different groups [8, 15, 26, 34]. In an animal model, Driessen showed that rigidified erythrocytes slow down the capillary blood flow (up to stasis) only if the blood pressure was lowered below the normotensive state [9]. On the other hand in extreme cases of sickle cell crisis, a dramatic near zero flow in cutaneous capillaries can occur [21, 24] because of the extremely reduction of the deformability of the sickled erythrocytes [27] together with their adhesiveness [3].

The problem is more complicated in cases of a pathologically elevated erythrocyte aggregation. Under these conditions a massive sludging can be found - firstly described by M.H. Knisely's group in conjunctival vessels - even in very small cutaneous capillaries [12]. Big erythrocyte aggregates pass very slowly through the capillary while immediately thereafter a single cell perfusion with significantly higher velocities in the same capillary can occur. In diabetic patients it can be shown that the erythrocyte aggregation correlated well with the mean erythrocyte velocity in nailfold capillaries in diabetic patients in a big clinical study with 1256 subjects [18].

In general, rheological phenomena are probably capable of deteriorating microvascular perfusion. However, while the as yet unknown regulatory mechanisms of the perfusion are intact, a direct impact is limited. In most patients, microcirculatory disorders are associated with pathologically changed rheological parameters, both processes seemingly in correlation to the extent of the disease. Therefore it can be assumed that in the natural course of arterial diseases endothelial cells, blood cells and the plasma also undergo pathological changes (this can be shown at least in patients with hypertension, diabetes mellitus, and with arterial occlusive disease) [1, 17, 19, 22].

In the case of arterial disease there is a line of events firstly mentioned by A.M. Ehrly [11]: rheological changes could induce and enhance a circulus vitiosus: vessel wall changes (stenosis) - a drop in pressure - a decreased blood flow - an acidosis enhancing the rheological defects - a.s.o.

Sudden increases in blood or in plasma viscosity (in combination with erythrocyte aggregation), lead to a viscosity-dependent drop in capillary perfusion in patients with coronary artery disease. This could be shown in ipsilateral nailfold capillaries by comparison of the effects of injection of high viscous or lower viscous radiographic contrast media into the A. axillaris [32] or in double blind studies in which the injection of high viscous iodinated contrast media were compared with the injection of isotonic sodium chloride solutions [2, 20]. On the other hand, an increase in capillary perfusion could be found after HELP apheresis with a plasma viscosity decrease of more than 20% in patients with macroglobulinaemia [7] or in heart-transplanted patients with transplant vasculopathy [28].

This shows that acute falls in plasma viscosity – though a concomitant decrease in erythrocyte aggregation also occurs because the fibrinogen concentration decreases [23, 30] - induce significant and relevant changes of capillary perfusion in humans. Studies to show the effects of changes of single rheological parameters (erythrocyte deformability, erythrocyte aggregation or platelet aggregation) suffer from serious limitations, because in most cases the drugs or interventional therapies [5, 13, 14, 16] applied, induced a vasodilation or vasoconstriction in addition.

In conclusion it is known that plasma viscosity in acute *in vivo* studies and - but only in the extreme cases of sickle cell crisis - the erythrocyte deformability influences the capillary perfusion. There is evidence that pathologically elevated erythrocyte aggregation slows down the capillary perfusion as long as aggregates perfuse the capillary [18]. In first - unpublished studies - we could see that in cases of increased numbers of circulating platelet aggregates there was a decrease in capillary perfusion, which normalised after applying thrombocyte function inhibitors.

There is clearly more work to be done in this area and the role of the leukocyte has not even been mentioned.

References

- [6] Chien S. Rheology in the microcirculation in normal and low flow states. *Adv Shock Res.* 1982;8:71-80.
- [10] Ehrly AM. Improvement of flow properties of blood: a new principle for drug therapy of chronic peripheral arterial vascular disorders. *Vasa Suppl.* 1973;1:1-18.
- [29] Rampling MW. Hyperviscosity as a complication in a variety of disorders. *Semin Thromb Hemost.* 2003;29(5):459-65.
- [31] Schmid-Schönbein H. Blood rheology and physiology of microcirculation. *Ric Clin Lab.* 1981;11 Suppl 1:13-33.
- [18] Jung F, Mrowietz C, Hiebl B, Franke RP, Pindur G, Sternitzky R. Influence of rheological parameters on the velocity of erythrocytes passing nailfold capillaries in humans. *Clin Hemorheol Microcirc* 2011; 48(1): 129-39.
- [33] Studer A, Hunziker T, Lütolf O, Schmidli J, Chen D, Mahler F. Quantitative nailfold capillary microscopy in cutaneous and systemic lupus erythematosus and localized and systemic scleroderma. *J Am Acad Dermatol.* 1991;24(6 Pt 1):941-5.
- [4] Barras JP. Blood rheology - general review. *Bibl Haematol.* 1969;33:277-97.
- [25] Lipowsky HH. Blood rheology aspects of the microcirculation. In : OK Baskurt, MR Hardeman, MW Rampling , HJ Meiselman (eds). *Handbook of Hemorheology and Hemodynamics.* IOS Press,Amsterdam, 2007, 307-321
- [8] Drenckhahn D. Zellmotilität und zytoplasmatische Filamentsysteme in Gefäßendothelzellen. In: K. Messmer, F. Hammersen (Eds). *Struktur und Funktion endothelialer Zellen.* Karger Verlag, Basel, 1983; 60-79.
- [15] Jung F, Franke RP. Extreme reduction of the capillary lumen in segments of the venular legs of human cutaneous capillaries. *Microvasc Res.* 2009;78:241 – 245.

- [26] Lübbers DW, et al. Contractile properties of frog capillaries tested by electrical stimulation. *Bibl anat.* 1979;17:3-10.
- [34] Tyml K, Weigelt H, Lübbers DW. Occurrence of the "capillary contractility" phenomenon and its significance in the distribution of microvascular flow in frog skeletal muscle. *Microvasc Res.* 1984;27(2):135-51.
- [9] Driessen GK, Fischer TM, Haest CW, Inhoffen W, Schmid-Schönbein H, Flow behaviour of rigid red blood cells in the microcirculation. *Int J Microcirc Clin Exp.* 1984;3(2):197–210.
- [24] Kord Valeshabad A, Wanek J, Zelkha R, Lim JI, Camardo N, Gaynes B, Shahidi M. Conjunctival microvascular haemodynamics in sickle cell retinopathy. *Acta Ophthalmol.* 2015;93(4):e275-80.
- [21] Kaul DK, Finnegan E, Barabino GA. Sickle red cell-endothelium interactions. *Microcirculation.* 2009;16(1):97-111.
- [27] Mozar A, Connes P, Collins B, Hardy-Dessources MD, Romana M, Lemonne N, Bloch W, Grau M. Red blood cell nitric oxide synthase modulates red blood cell deformability in sickle cell anemia. *Clin Hemorheol Microcirc.* 64 (1),2016, 47-53.
- [3] Ballas SK, Connes P. The paradox of the serrated sickle erythrocyte: The importance of the red blood cell membrane topography. *Clin Hemorheol Microcirc.* 2015;16;63(2):149-52.
- [12] Frayser R, Knisely WH, Barnes R, Satterwhite WM Jr. In vivo observations on the conjunctival circulation in elderly subjects. *J Gerontol.* 1964;19:494-500.
- [1] Arend O, Wolf S, Jung F, Bertram B, Pöstgens H, Toonen H, Reim M. Retinal microcirculation in patients with diabetes mellitus: dynamic and morphological analysis of perifoveal capillary network. *Brit J Ophthalmol* 1991; 75: 514 - 518.
- [19] Jung F, Pindur G, Ohlmann P, Spitzer G, Sternitzky R, Franke RP, Leithäuser B, Wolf S, Park JW. Microcirculation in hypertensive patients. *Biorheology* 2013; 50(5): 241-55.
- [17] Jung F, Krüger A, Pindur G, Sternitzky R, Franke RP, Gori T. Tissue oxygen partial pressure in the tibialis anterior muscle in patients with claudication before, during and after a two-stage treadmill stress test. *Biorheology.* 2014;51(2-3):207-14.
- [22] Kiesewetter H, Jung F, Ladwig KH, Waterloh E, Roebuck P, Schneider R, Kotitschke G, Bach R. Predictor function of hemorheologic parameters with reference to the incidence of manifest circulatory disorders: Concept of the Aachen study. *Klin Wochenschr.* 1986;64(14):653-62
- [11] Ehrly AM. New pathophysiological concept of ischemic diseases: Microcirculatory blood maldistribution. *Bibl Anat.* 1983;20:456-460.
- [32] Spitzer S, Münster W, Sternitzky R, Bach R, Jung F. Influence of Iodixanol-270 and Iopentol-150 on the microcirculation in man: influence

- of viscosity on capillary perfusion. *Clin Hemorheol Microcirc* 1999; 20: 49 - 55.
- [2] Bach R, Jung F, Scheller B, Hummel B, Özbek C, Spitzer S, Schieffer H. Influence of a non-ionic radiography contrast medium on the microcirculation. *Acta Radiologica*. 1996; 37: 214 - 217.
- [20] Jung F, Spitzer S, Mrowietz C, Sternitzky R, Bach R. Influence of radiographic contrast media (Iomeprol350 versus Iopentol350) on cutaneous microcirculation: single-center prospective randomized double-blind phase IV study in parallel group design. *J Invas Cardiol*. 1999; 11: 2 - 8.
- [7] Dobberstein H, Solbach U, Weinberger A, Wolf S. Correlation between retinal microcirculation and blood viscosity in patients with hyperviscosity syndrome. *Clin Hemorheol Microcirc*. 1999;20(1):31-5.
- [28] Park JW, Mrowietz C, Schüler S, Labarrere C, Jung F. Cutaneous microcirculation in cardiac allograft recipients with severe hypercholesterolemia before, during, and after the first HELP apheresis. *Applied Cardiopulmonary Pathophysiology* 2000; 9: 19-25.
- [23] Kirschkamp T, Schmid-Schönbein H, Weinberger A, Smeets R. Effects of fibrinogen and alpha2-macroglobulin and their apheretic elimination on general blood rheology and rheological characteristics of red blood cell aggregates. *Ther Apher Dial*. 2008 Oct;12(5):360-7.
- [30] Rampling MW, Whittingstall P. A comparison of five methods for estimating red cell aggregation. *Klin Wochenschr*. 1986;64(20):1084-8.
- [13] Gori T, Münzel T. Endothelial dysfunction after stenting and scaffolding of coronary arteries. *Clin Hemorheol Microcirc*. 2014;58(1):175-81.
- [5] Bilsborough W, O'Driscoll G, Stanton K, Weerasooriya R, Dembo L, Taylor R, Green D. Effect of lowering tumour necrosis factor-alpha on vascular endothelial function in Type II diabetes. *Clin Sci (Lond)*. 2002 Aug;103(2):163-9.
- [14] Grossmann M, Braune J, Ebert U, Kirch W. Dilatory effects of phosphodiesterase inhibitors on human hand veins in vivo. *Eur J Clin Pharmacol*. 1998 Mar;54(1):35-9.
- [16] Jung F, Kiesewetter H, Mrowietz C, Leipnitz G, Braun B, Wappler M, Scheffler P, Wenzel E. Hemorrheological, micro- and macrocirculatory effects of naftidrofuryl in an acute study: a randomized, placebo-controlled, double-blind individual comparison. *Int J Clin Pharmacol Ther Toxicol*. 1987;25(9):507-14.